

2/7/00



CHOATE, HALL & STEWART

A PARTNERSHIP INCLUDING PROFESSIONAL CORPORATIONS

EXCHANGE PLACE


53 STATE STREET

BOSTON, MASSACHUSETTS 02109-2891

TELEPHONE (617) 248-5000

FACSIMILE (617) 248-4000

EXPRESS MAIL NO. EK152416823US



February 4, 2000

Assistant Commissioner for Patents
Washington, DC 20231
Box Patent Application

TRANSMITTAL LETTER

Dear Sir:

Please find enclosed a patent application as follows:

Applicant(s): Burt D. Ensley

Title: Wound Healing Compositions and Methods Using Tropoelastin and Lysyl Oxidase

No. Pages Specification: 27; No. Pages Claims: 4; No. Pages of Drawings: 1 ; No. Pages Abstract: 1;

Enclosed is: Unexecuted Declaration; Unexecuted Appointment of Attorney; Unexecuted Verified Statement (Declaration) Claiming Small Entity Status 37 CFR 1.9(f) and 1.27(b) Independent Inventor; Check for \$429.00; and Return Postcard.

Basic Fee: \$345.00

Additional Fees:

Total Number of Claims in excess of 20: $25 - 20 = 5 \times \$9$ \$45.00

Number of independent claims minus 3 times: $4 - 3 = 1 \times \$39$ \$39.00


Total Filing Fee \$429.00

Please charge any additional fees that are required or credit any overpayments to our Deposit Account No. 03-1721.

If this application is found otherwise to be INCOMPLETE, or if at any time it appears that a TELEPHONE CONFERENCE with counsel would helpfully advance prosecution, please telephone the undersigned at any time.

Kindly acknowledge receipt of the foregoing application by returning the self-addressed postcard.

Respectfully submitted,


Brenda H. Jarrell, Ph.D., Reg. No. 39,223

ATTORNEY DOCKET NO.: 2001605-0007

Applicant: Burt D. Ensley
Serial No.:
Filed: February 4, 2000
For: WOUND HEALING COMPOSITIONS AND METHODS USING
TROPOELASTIN AND LYSYL OXIDASE

VERIFIED STATEMENT (DECLARATION) CLAIMING SMALL ENTITY STATUS
37 CFR 1.9(f) and 1.27(b)
INDEPENDENT INVENTOR

As a below-named inventor, I hereby declare that I qualify as an independent inventor as defined in 37 CFR 1.9(c) for purposes of paying reduced fees under section 41(a) and (b) of Title 35, United States Code with regard to the invention titled:

TITLE: Wound Healing Compositions and Methods Using Tropoelastin and Lysyl Oxidase

by:

INVENTOR: Burt D. Ensley

described in:

- ☒ (X) the specification filed herewith
- ☐ () U.S. Patent Application Serial Number
filed
- ☐ () U.S. Patent Number _____
issued _____

I have not assigned, granted, conveyed, or licensed and am under no obligation under contract or law to assign, grant, convey, or license, any rights in the invention to any person who could not be classified as an independent inventor under 37 CFR 1.9(c) if that person had made the invention, or to any concern that would not qualify as a small business concern under 37 CFR 1.9(d) or a non-profit organization under 37 CFR 1.9(e).

Each person, concern, or organization to which I have assigned, granted, conveyed, or licensed, or am under an obligation under contract or law to assign, grant, convey, or license any rights in the invention is listed below:

- ☒ (x) no such person, concern, or organization
- ☐ () persons, concerns, or organizations listed below*:

*NOTE: Separate verified statements are required from each named person, concern, or organization having rights to the invention, averring to their status as small entities. (37 CFR 1.27).

I acknowledge the duty to file, in this application or patent, notification of any change in status resulting in loss of entitlement to small entity status prior to paying, or at the time of paying, the earliest of the issue fee or any maintenance fee due after the date on which status as a small entity is no longer appropriate. (37 CFR 1.28(b))

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under §1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application, any patent issuing thereon, or any patent to which this verified statement is directed.

NAME OF INVENTOR:

Burt D. Ensley

ADDRESS OF INVENTOR:

7 Colts Neck Drive, Newtown, PA 18940

SIGNATURE: _____ DATE: _____

Exchange.3090236.1

WOUND HEALING COMPOSITIONS AND METHODS USING TROPOELASTIN AND LYSYL OXIDASE

Background of the Invention

- 5 Wound healing is a complex biological process that involves many different cell types, many different cytokines, the extracellular matrix (ECM), and numerous interactions among them. Most wounds heal rapidly and efficiently within a week or two; however, the result is neither aesthetically nor functionally perfect. Wound contraction and scar formation are currently unavoidable results of wound healing.
- 10 Scar tissue is less flexible than normal skin and can be cosmetically disfiguring, and wound contraction can lead to joint disablement (Lamme *et al.*, *J. Histochem. Cytochem.* 44:1311, 1996). Scars lack elastin and consist of poorly reconstituted collagen matrix in dense parallel bundles rather than the mechanically efficient basket-weave meshwork of collagen in unwounded dermis (Martin, *Science* 276:75, 1997).
- 15 Two major goals of wound-healing biology are more rapid wound healing and more perfect reconstruction of the damaged parts. Compositions and methods useful in accomplishing these goals are currently needed.

Wound Healing

- 20 Wound healing has been divided into a number of overlapping phases. These include fibrin clot formation, recruitment of inflammatory cells, reepithelialization, and matrix formation and remodeling. Immediately after tissue injury, blood vessel disruption leads to the extravasation of blood and concomitant platelet aggregation and

blood coagulation resulting in fibrin clot formation. Activated platelets trapped within the fibrin clot degranulate and release a variety of cytokines and growth hormones. These cytokines and growth hormones help to recruit inflammatory cells to the site of injury, to stimulate angiogenesis, and to initiate the tissue movements associated with

5 reepithelialization and connective tissue contraction.

Neutrophils and monocytes are recruited to the site of injury by a number of chemotactic signals including the growth factors and cytokines released by the degranulating platelets, formyl methionyl peptides cleaved from bacterial proteins, and the by-products of proteolysis of fibrin and other matrix proteins. Neutrophil

10 infiltration ceases after a few days, but macrophages continue to accumulate by continued recruitment of monocytes to the wound site. Activated macrophages release growth factors and cytokines thereby amplifying the earlier signals from the degranulating platelets.

Formation of granulation tissue and reepithelialization of the wound site begins

15 after several hours (Clark, *J. Am. Acad. Dermatol.* 13:701, 1985).

Reepithelialization is performed by the basal keratinocytes which lose their attachments to the basal lamina and crawl over the provisional matrix of fibrin and fibronectin, and underlying matrix. Some hours after the onset of migration, epidermal cells begin to reproduce and thereby provide the cells needed to replace

20 those lost during the injury. Keratinocyte proliferation is regulated by keratinocyte growth factor and members of the epidermal growth factor (EGF) family. In order to migrate through the fibrin clot, the keratinocytes must dissolve the fibrin barrier in

front of them. Plasmin is the chief fibrinolytic enzyme used in this process.

Reepithelialization is made easier by the underlying contractile connective tissue, which shrinks to bring the wound margins toward one another. Epidermal migration ceases when the wound surface has been covered by a monolayer of cells.

5 Cells of the new epidermis undergo the standard differentiation program of cells in the outer layers of unwounded epidermis. A new stratified epidermis is, thereby, reestablished from the margins of the wound inward. Matrix formation and remodeling begins simultaneously with reepithelialization. The matrix is constantly altered over the next several months with the elimination of the fibronectin from the
10 matrix and the accumulation of collagen that provides the residual scar with increasing tensile strength. Elastin fibers, which are responsible for the elasticity of tissue, are only detected in human scars years after the injury (Compton *et al.*, *Lab. Invest.* 60:600, 1989).

15 *Methods to Promote Wound Healing*

 In the past, many methods have been proposed and tested to promote wound healing and limit scarring; however, better methods and compositions are still needed. These older methods include cyanoacrylate tissue adhesives, a combination of epidermal transplantation and a collagen/elastin dermal substitute, application of
20 collagen and glycosaminoglycans to the site of injury, and biocompatible adhesives with collagen.

One of the more popular methods is the use of cyanoacrylate tissue adhesives. These adhesive have been used in place of and in conjunction with sutures. Cyanoacrylate adhesives have been used in cases ranging from cardiac surgery (Robicsek *et al.*, *J. Card. Surg.* 9:353, 1994) to simple lacerations in the pediatric
5 population (Penoff, *Plast. Reconstr. Surg.* 103:730, 1999). One study has found that cyanoacrylate tissue adhesive may be the preferred method in terms of cosmetic appearance for the cutaneous closure of facial lacerations oriented against Langer's lines (Simon *et al.*, *J. Emerg. Med.* 16:185, 1998).

In a study of the healing of full-thickness wounds in pigs, a dermal matrix
10 consisting of native bovine collagen coated with elastin hydrolysate was found to serve as a template for dermal tissue regeneration in combination with an epidermal transplantation. This combination treatment reduced wound contraction and improved tissue regeneration (Lamme *et al.*, *J. Histochem. Cytochem.* 44:1311, 1996).

U.S. Patent No. 4,837,024, issued June 6, 1989, to Michaeli *et al.*, discloses
15 an article to promote healing of a surface wound. A suspension of particles of collagen and a glycosaminoglycan is contacted with the wound surface. Collagen is a major component of the ECM and helps to promote wound healing. The glycosaminoglycan is chemotactic of fibroblasts and/or endothelial cells. The collagen/glycosaminoglycan is applied to the wound and maintained in contact with
20 the wound for an extended period of time, *i.e.*, during the entire healing process or until at least closure of the wound by new tissue. The application promotes the

vascularization of the wound, attracts fibroblasts and endothelial cells, and generally provides a favorable environment for the cells during the healing process.

U.S. Patent No. 5,614,587, issued March 25, 1997, to Rhee *et al.*, and U.S. Patent No. 5,744,545, issued April 28, 1998, to Rhee *et al.*, disclose a composition
5 suitable for use as a bioadhesive and a method for using such a composition. The composition comprises fibrillar collagen, a fiber disassembly agent, and a multifunctionally activated synthetic hydrophilic polymer. The collagen and polymer are mixed to initiate cross-linking, the collagen-polymer mixture is then applied to a first surface before substantial cross-linking has occurred, and then a second surface is
10 brought in contact with the first surface. The composition is optically clear so that it could be used in ophthalmic applications, and the composition comprises biocompatible, non-immunogenic components which leave no toxic, potentially inflammatory or immunogenic reaction products at the tissue site of administration.

15 An invention which would promote healing and lessen scarring would be of great value.

Summary of the Invention

The present invention provides compositions and methods useful in the
20 promotion of wound healing. These compositions comprise virgin monomers of tropoelastin and the cross-linking enzyme, lysyl oxidase. The method comprises mixing these two components of the composition together and applying them to a

wound before substantial cross-linking has occurred. The tropoelastin monomers and lysyl oxidase only come in contact with each other immediately before application to the wound or during application to the wound.

Without wishing to be bound by any particular theory, we propose that the

5 lysyl oxidase catalyzes the oxidative deamination of the lysine residues of the tropoelastin monomers at the site of the wound. Then in a non-enzymatic step, cross-links form between the tropoelastin monomers as well as between tropoelastin monomers and other proteins of the extracellular matrix such as collagen. The cross-linked elastin at the site of injury helps to hold the injured tissue together and thereby

10 promotes healing. The elastin is also chemotactic for fibroblasts, endothelial cells, and inflammatory cells, thereby promoting healing in another manner. Elastin at the site of injury also helps to lessen scarring since scar tissue is devoid of elastin, and elastin is an important component of uninjured skin. The cross-linked elastin also generally provides a favorable environment for the cells that participate in the healing

15 process.

In preferred embodiments of the invention, one or both of the tropoelastin monomers and lysyl oxidase are made recombinantly and purified to homogeneity using standard techniques. The purified tropoelastin and lysyl oxidase may then be suspended in a liquid, such as an aqueous solution (*e.g.*, water or saline) or an

20 organic solvent, or provided in a dry powder form, or in a lyophilized form. These two components, tropoelastin and lysyl oxidase, are kept separate from each other until right before use. In another embodiment, the lysyl oxidase is kept in an inactive

form in the presence of the tropoelastin, and the lysyl oxidase not activated until right before use. In yet another embodiment, the two components are applied to the wound separately.

In the method of the present invention, the composition may be applied only
5 once at the time of the injury or more than once over the course of wound healing. The composition of tropoelastin and lysyl oxidase may also be used in conjunction with sutures, staples, or adhesive strips in closing the wound.

The composition of the present invention may also be used in promoting the healing of wounds involving structures with elastic fibers such as arteries, lung tissue,
10 or skin. In particular, the composition may be used in surgeries involving arteries, lungs, or the skin.

In another embodiment of the present invention, the tropoelastin comprises only portions of the tropoelastin protein, preferably containing at least one cross-linking domain. In another embodiment, the lysyl oxidase comprises only an active
15 portion of the enzyme.

The preferred compositions of the present invention are biocompatible, non-toxic, and non-immunogenic, and potentially inflammatory or immunogenic reaction products at the site of administration are avoided.

20

Description of the Drawings

Figure 1 shows the reaction catalyzed by lysyl oxidase, the oxidative deamination of lysine. A variety of non-enzymatic cross-linking reactions and their products are also shown.

5 Definitions

Animal refers to human as well as non-human animals. Non-human animals include mammals, birds, reptiles, amphibians, and fish. Preferably, the non-human animal is a mammal (*i.e.*, a rodent, a mouse, a rat, a rabbit, a monkey, a dog, a cat, or a pig). The term includes transgenic animals.

10 *Biocompatible* refers to material that is not toxic to the body, is not carcinogenic, and does not induce inflammation in body tissues.

Biodegradable refers to material that is degraded by normal bodily processes resulting in products which are readily disposable by the body and do not accumulate in the body.

15 *Elastin* refers to the cross-linked extracellular matrix protein. *In vivo*, tropoelastin monomers are cross-linked by lysyl oxidase to form elastin.

Isolated refers to a protein being substantially purified away from contaminating matter. In a preferred embodiment, the protein is greater than 70% pure, greater than 80% pure, or greater than 90% pure. In a particularly preferred
20 embodiment, the protein is greater than 95% pure or 98% pure.

Lysyl oxidase refers to a catalyst which is able to catalyze the oxidative deamination of the amino acid lysine as shown in Figure 1. In preferred

embodiments of the invention, the catalyst is a lysyl oxidase protein enzyme. The genes encoding such enzymes have been cloned from a variety of organisms (Hämäläinen *et al.*, *Genomics* 11:508, 1991; Trackman *et al.*, *Biochemistry* 29:4863, 1990; incorporated herein by reference). In accordance with the present invention,

5 the lysyl oxidase employed is preferably selected to match as closely as possible the individual subject to which the compositions are to be applied. For example, human lysyl oxidase, which has been cloned and characterized (Hämäläinen *et al.* "Molecular Cloning of Human Lysyl Oxidase and Assignment of the Gene to Chromosome 5q23.3-31.2" *Genomics* 11(3):508-516, 1991; incorporated herein by reference) is

10 preferably utilized in the treatment of wounds involving human tissue. However, in other embodiments the lysyl oxidase may be obtained from any species. When compared to the lysyl oxidase protein of the target species, useful analogs of lysyl oxidase generally exhibit at least 60% homology, preferably at least about 70% homology, more preferably at least about 80% homology, and most preferably at least

15 90%, 95%, or 99% homology, with a segment of 20 amino acid residues, preferably with more than 40 amino acid residues, more preferably yet with substantially the entire sequence of the target species lysyl oxidase. For compositions to be applied to humans, it is particularly preferred that the lysyl oxidase shows homology to a segment from residues 153-417 and residues 201-417 of the sequence of human lysyl

20 oxidase.

The lysyl oxidase employed in the practice of the present invention may be modified either chemically or genetically *in vivo* or *in vitro*. Examples of non-

sequence chemical modifications include phosphorylation, acetylation, methylation, carboxylation, hydroxylation, and glycosylation. Examples of genetic modifications include changes in the amino acid sequence, truncation of the polypeptide chain, addition of at least one amino acid, and deletion of at least one amino acid.

- 5 Techniques for producing such chemical and/or genetic modifications are well known in the art. Any modified protein or polypeptide that retains the ability to catalyze the oxidative deamination of lysine on tropoelastin is useful in the practice of the present invention.

- Preferred analogs of lysyl oxidase include human lysyl oxidase or biologically
10 active fragments thereof, whose sequence differ from the wild type by one or more conservative amino acid substitutions or by one or more non-conservative amino acid substitutions, deletions, or insertions which do not abolish lysyl oxidase biological activity. Conservative substitutions typically include the substitution of one amino acid for another with similar characteristics (*i.e.*, substitutions within the following
15 groups: valine, glycine; glycine, alanine; valine, isoleucine, leucine; aspartic acid, glutamic acid; asparagine, glutamine; serine, threonine; lysine, arginine; and phenylalanine, tyrosine). Other conservative substitutions are known by those skilled in the art.

- In still other embodiments of the invention, lysyl oxidase may be a non-
20 polypeptide catalyst with the ability to catalyze the oxidative deamination of lysine.

Protein, peptide, or polypeptide refers to a polymer of amino acids, and these terms are used interchangeably. The polymer may include natural or unnatural amino

acids. The protein or polypeptide may be produced *in vitro* or *in vivo* via natural, recombinant, synthetic, or other means. The protein or polypeptide may have post-translational modifications or may have been modified chemically to include phosphorylation, glycosylation, farnesylation, acetylation, methylation, oxidation of
5 thiols, *etc.*

Recombinant can refer to organisms, cells, nucleic acids, and proteins.

Recombinant cells and organisms are cells and organisms containing recombinant DNA.

Recombinant DNA refers to a nucleic acid sequence which is not normally found in nature. Usually this term refers to two or more pieces of DNA spliced together to form

10 an unnatural product. Recombinant protein is protein produced from recombinant DNA (*i.e.*, a nucleic acid which differs from that which occurs in nature). In producing a recombinant protein, the regulatory sequences of the gene encoding the protein are usually different than the ones that occur in the natural gene. The gene may also have been placed in an organism which normally does not possess the gene in order to produce
15 that protein in the desired organism.

Target species refers to the species on which the composition will be applied. A target species can be any animal including humans. Preferably, the target species is mammalian, more preferably a domestic mammal (*e.g.*, dog, cat, cow, horse, rabbit, goat, hamster) or a rodent (*e.g.*, rat, mouse), and most preferably human.

20 *Tropoelastin* refers to monomer polypeptides which, when cross-linked, form elastin. The genes encoding tropoelastin have been cloned from a variety of organisms (Bressan *et al.* "Repeating structure of chick tropoelastin revealed by

complementary DNA cloning" *Biochemistry* 26:1497-1503, 1987; Raju *et al.*

"Primary structures of bovine elastin a, b, and c deduced from the sequences of cDNA clones" *J. Biol. Chem.* 262:5755-5762, 1987; Indik *et al.* "Alternative splicing of human elastin mRNA indicated by sequence analysis of cloned genomic and

5 complementary DNA" *Proc. Natl. Acad. Sci. USA* 84:5680-5684, 1987; Yeh *et al.*

"Structure of the bovine elastin gene and S1 nuclease analysis of alternative splicing of elastin mRNA in the bovine nuchal ligament" *Biochemistry* 28:2365-2370, 1989;

Pierce *et al.* "Heterogeneity of rat tropoelastin mRNA revealed by cDNA cloning" *Biochemistry* 29:9677-9683, 1990; each of which is incorporated herein by reference).

10 In accordance with the present invention, the lysyl oxidase employed is preferably selected to match as closely as possible the individual subject to which the compositions are to be applied. In preferred embodiments of the current invention, the human tropoelastin is utilized in the treatment of a wound involving human tissue. A variety of different isoforms of human tropoelastin are produced in nature (by
15 alternative splicing); any such isoform, or collection of isoforms, may be utilized in the practice of the present invention. In other embodiments, non-human tropoelastin may be employed; however, it is generally desirable to match the species of animal being treated to the species of the tropoelastin being used. The tropoelastin protein may be modified, as compared with naturally occurring human tropoelastin protein,
20 either chemically or genetically *in vivo* or *in vitro*. When compared to the tropoelastin protein of the target species, useful analogs of tropoelastin generally exhibit at least 60% homology, preferably at least about 70% homology, more

preferably at least about 80% homology, and most preferably at least 90%, 95%, or 99% homolgy, with a segment of 20 amino acid residues, preferably with more than 40 amino acid residues, more preferably yet with substantially the entire sequence of human tropoelastin. Examples of non-sequence chemical modifications include

- 5 phosphorylation, acetylation, methylation, carboxylation, hydroxylation, and glycosylation. Examples of genetic modifications would be changes in the amino acid sequence, truncation of the polypeptide chain, addition of at least one amino acid, and deletion of at least one amino acid. Techniques for producing such chemical and/or genetic modifications are well known in the art. Tropoelastin can be prepared
- 10 recombinantly or by chemical synthesis, but it cannot be purified from natural sources because it will have already been crosslinked.

Wild type refers to a nucleic acid sequence or amino acid sequence which is found in nature and has not been mutated. In a preferred embodiment, the wild type sequence is one of the more common sequences for a particular gene or protein found

15 in a particular species. In some instances, there may be several wild type sequences for a protein or gene for one particular species due to multiple alleles of a gene or multiple isoforms of a protein.

Wound refers to damaged biological tissue in the most general sense. In a preferred embodiment, the wound is a laceration of the skin. In other embodiments,

20 the wound may be an abrasion of the skin without two separated parts of tissue which need to be brought together. In still other embodiments, the wound may refer to a

surgical incision. In other preferred embodiments, the wound may involve damage to lung tissue, arterial walls, or other organs with elastic fibers.

Detailed Description of Preferred Embodiments

5 As described above, the present invention provides compositions and methods for promoting wound healing. These compositions comprise tropoelastin and lysyl oxidase. The method of the present invention involves contacting these two components together and applying the mixture to a wound before substantial cross-linking of the tropoelastin has taken place. In another embodiment, the tropoelastin
10 and lysyl oxidase are applied separately to the wound. A key aspect of the invention is that the tropoelastin must be applied to the wound before substantial cross-linking of the tropoelastin has occurred. To those skilled in the art, a variety of ways of applying the two substances wherein the tropoelastin is not substantially cross-linked will be clear. With the cross-linking reaction taking place at the wound site, cross-
15 links will form that will hold the tissue together. Also, the new formation of elastin will attract fibroblasts, inflammatory cells, and endothelial cell by chemotaxis, will result in less scarring, and will provide a suitable environment for the cells involved in the healing process.

20 *Elastin*

Elastic fibers in the extracellular space of the lungs, dermis, and large blood vessels contribute to the elasticity and resilience of these tissues. These elastic fibers

may comprise only a small (2-4%) portion of the dry weight of skin but may
comprise greater than 50% of the dry weight of large arteries (Rosenbloom *et al.*,
FASEB J. 7:1208, 1993). These fibers are composed of amorphous elastin and
microfibrils. Microfibrils are a complex of glycoproteins organized as small, 10- to
5 12-nm-diameter fibrils and serve as the scaffold onto which elastin is assembled.
Elastin is the major component (>90%) of the mature elastic fibers and has been well
characterized. The elastin protein is initially synthesized as tropoelastin monomers,
soluble polypeptides of *ca.* 72 kDa (Parks *et al.*, *Am. J. Respir. Cell Mol. Biol.*
2:399, 1990).

- 10 The tropoelastin gene exists as a single copy in the genome. The gene has
been isolated from several species, including human, bovine, chick, and rat (Bressan
et al. "Repeating structure of chick tropoelastin revealed by complementary DNA
cloning" *Biochemistry* 26:1497-1503, 1987; Raju *et al.* "Primary structures of bovine
elastin a, b, and c deduced from the sequences of cDNA clones" *J. Biol. Chem.*
15 262:5755-5762, 1987; Indik *et al.* "Alternative splicing of human elastin mRNA
indicated by sequence analysis of cloned genomic and complementary DNA" *Proc.*
Natl. Acad. Sci. USA 84:5680-5684, 1987; Yeh *et al.* "Structure of the bovine elastin
gene and S1 nuclease analysis of alternative splicing of elastin mRNA in the bovine
nuchal ligament" *Biochemistry* 28:2365-2370, 1989; Pierce *et al.* "Heterogeneity of
20 rat tropoelastin mRNA revealed by cDNA cloning" *Biochemistry* 29:9677-9683, 1990;
each of which is incorporated herein by reference); the entire bovine and human
elastin genes have been isolated and sequenced (Indik *et al.*, *Proc. Natl. Acad. Sci.*

USA 84:5680, 1987; Yeh *et al.*, *Biochemistry* 28:2365, 1989; Bashir *et al.*, *J. Biol. Chem.* 264:8887, 1989; each of which is incorporated herein by reference).

Elucidation of the amino acid sequence indicates alternating segments of cross-link domains and hydrophobic domains. The cross-link domains are characterized by the presence of lysine residues separated by two to three alanine residues, and the hydrophobic domains are found to be rich in hydrophobic residues such as glycine, alanine, valine, and proline. This composition results in hydrophobic interactions which are thought to be responsible for the elasticity of the fibers. The cross-link and hydrophobic domains are encoded by separate exons of the gene.

Extensive alternative splicing of the primary tropoelastin transcript yields the diversity of tropoelastin isoforms seen at the protein level. The human tropoelastin gene consists of 34 separate exons spanning a total of *ca.* 45 kB of genomic DNA. Six of these exons have been reported to be subject to alternative splicing (Uitto *et al.*, *Biochem. Soc. Transact.* 19:824, 1991).

Newly synthesized pre-tropoelastin (containing the signal peptide) undergoes intracellular post-translational modifications including hydroxylation of certain proline residues to form 4-hydroxyproline and removal of the signal peptide. The tropoelastin monomers are then secreted into the extracellular milieu where they assemble into functional fibers (fibrillogenesis) and are cross-linked to form insoluble elastin (Uitto *et al.*, *Biochem. Soc. Transact.* 19:824-829, 1991).

The assembly of elastic fibers (fibrillogenesis) takes place at unique sites close to the cell membrane, generally in infoldings of the cell surface. Microfibrils are the

first component to appear and are found to be grouped in small bundles near the plasma membrane. Elastin then appears as amorphous material in discrete loci within each microfibrillar bundle. The microfibrils are thought to align the tropoelastin molecules so that the cross-linking regions are juxtaposed. Lysyl oxidase then

5 oxidizes the terminal amino groups on the side chains of the lysine residues of the cross-linking regions. The oxidized side chains then undergo non-enzymatic condensation reactions to form the cross-links. The notion that tropoelastin monomers are secreted from the cell and diffuse onto the surface of growing fibers seems to be inadequate to explain the efficiency of the whole process and the variable
10 forms of elastic fibers in different tissues. Rather, increasing evidence has supported the idea that helper proteins are needed in the secretion and fiber assembly steps. Given that tropoelastin monomers produced in organisms are rapidly incorporated into insoluble elastin fibers, purification of tropoelastin from anything other than recombinant sources is not practical. Thus, in accordance with the present invention,
15 virgin tropoelastin monomers may be provided by any available method, including, for example, chemical synthesis and standard recombinant techniques known in the art.

In one preferred embodiment, the tropoelastin monomers are synthesized using available chemical synthetic methods. For example, tropoelastin monomers can be
20 synthesized using an appropriate solid state synthetic procedure (Steward *et al.*, *Solid Phase Synthesis*, Freeman, San Francisco, CA, 1968). A preferred method is the Merrifield process (Merrifield, *Recent Prog. Hormone Res.* 23:451, 1967).

Alternatively, tropoelastin monomers may be prepared by recombinant techniques. Techniques for the overexpression and purification of recombinant proteins in a wide variety of cell types are well known in the art ("Gene Expression Technology," *Methods in Enzymology*, vol. 185 (D. V. Goeddel, ed.), Academic Press Inc., 1990; incorporated herein by reference). Any such techniques can be employed in accordance with the present invention. Since tropoelastin monomers are normally secreted from the cell, a preferred way to produce the monomers for the present invention would be to introduce the gene with the signal sequence intact and under the control of a strong promoter into a cell and grow the recombinant cells in culture. Preferably, the recombinant cells are eukaryotic (*i.e.*, *Saccharomyces cerevisiae*, *Pichia pastoris*), more preferably the cells are of mammalian origin (*i.e.*, COS cell line, CHO cell line), and most preferably the cells are of a human origin. In certain preferred embodiments of the invention, the recombinant cells are derived from human fibroblasts. In another preferred embodiment, the recombinant cells are recombinant bacteria (*e.g.*, *E. coli*). The desired tropoelastin monomers would be secreted into the media, and the protein could be purified to homogeneity from the media. To improve the ease at which the protein could be purified, the cells expressing the tropoelastin would be grown in serum-free media, thereby minimizing the amount of protein in the media to start with. Production of the tropoelastin monomers in mammalian cells such as fibroblasts would insure correct folding and post-translational modification of the protein. This could be important since some of the proline residues in tropoelastin are hydroxylated posttranslationally.

Once the recombinant tropoelastin of the present invention is expressed, the protein may be isolated and purified by standard methods including chromatography (e.g., ion exchange, affinity, size exclusion, and hydroxyapatite chromatography), centrifugation, or differential solubility, or by any other available technique for the

5 purification of proteins (Scopes, *Protein Purification Principles and Practice 2nd Edition*, Springer-Verlag, New York, 1987; incorporated herein by reference). For immunoaffinity chromatography in particular, the protein may be isolated by binding it to an affinity column comprising antibodies that were raised against that protein and were affixed to a stationary support. Alternatively, affinity tags such as an influenza

10 coat sequence, poly-histidine, or glutathione-S-transferase can be attached to the protein by standard recombinant techniques to allow for easy purification by passage over the appropriate affinity column.

Any method of producing tropoelastin known in the art can be used as long as the tropoelastin is able to be cross-linked to form elastin and does not initiate negative

15 (e.g., immune reactions) reactions once introduced into or applied onto the recipient's body.

If particular isoforms of tropoelastin are needed, the appropriate cDNA(s) can be introduced into the cells for overexpression. Different isoforms of tropoelastin are found in different individuals within a species, in different locations within one

20 individual, and in different tissues within an individual. In preferred embodiments of the present invention, a certain isoform of tropoelastin or mixture of isoforms is used for a particular individual being treated, a particular type of wound, or a particular

type of tissue. For example, it may be desirable to provide a composition comprising multiple different tropoelastin isoforms, whose identity and/or relative quantity are selected to match as closely as possible the profile of isoforms naturally produced by the individual to whom the composition is being applied, at the site of application. If

5 the natural isoform distribution in that individual (or a representative thereof, *e.g.*, another member, or an averaged collection of members, of the same species) and/or at that site, is not known in advance it may readily be determined by one of ordinary skill in the art using standard techniques such as, for example, gene sequencing, restriction fragment length polymorphisms, Western blotting, immunoassay, Northern

10 blotting, Southern Blotting, isoelectric focusing, SDS-polyacrylamide gel electrophoresis, *etc.* In a preferred embodiment, the individual isoforms of tropoelastin might be prepared separately and mixed at specific ratios later to simulate the ratio of isoforms in a specific tissue of an individual. The use of tropoelastin isoforms native to the tissue would minimize any immune reactions.

15 In another preferred embodiment of the invention, a modified version of tropoelastin is provided which is not susceptible to cross-linking. The tropoelastin may not be susceptible to cross-linking for a number of reasons including, for example, its binding another protein or peptide, or its not being folded correctly. In order for the tropoelastin to be cross-linked, the binding protein or peptide would

20 need to be removed, or the tropoelastin would need to be refolded. The binding peptide or protein may be removed by competing off the peptide or protein; cleavage of the binding peptide or protein; conformational change of one of the proteins; *etc.*

If the tropoelastin is not folded correctly, a protein from the family of chaperonins (*e.g.*, GroEL, GroES) or heat shock proteins (*e.g.*, Hsp60, Hsp40, Hsp70) may be added to re-fold the tropoelastin so that it may be cross-linked. In this embodiment, the virgin tropoelastin monomers and modified lysyl oxidase can be stored together
5 without the risk of cross-linking the tropoelastin monomers.

Lysyl Oxidase

The high degree of cross-linking found in the elastic fibers contributes to their proper function. Lysyl oxidase is the enzyme that catalyzes the oxidative deamination
10 of lysine residues leading to the non-enzymatic condensation of the modified lysine side chains. This same enzyme is involved in collagen cross-link formation as well. All but about 5 of the 34 lysine residues of tropoelastin participate in some form of cross-link resulting in a highly insoluble polymer (Rosenbloom *et al.*, *FASEB J.* 7:1208, 1993).

15 Lysyl oxidase is an extracellular, copper-requiring enzyme which has been purified to homogeneity from several animal sources and found to have a molecular weight of *ca.* 30 kDa (Stassen, *Biochim. Biophys. Acta* 438:49, 1976; Siegel, *Int. Rev. Connect. Tissue Res.* 8:73, 1979; Sullivan *et al.*, *J. Biol. Chem.* 257:13520, 1982; Kagan, *Biology of Extracellular Matrix* 1:321, 1986; Kuivaniemi *et al.*, *J. Biol.*
20 *Chem.* 259:6996, 1984). The genes encoding lysyl oxidase have been cloned from human and rat cDNA libraries (Hämäläinen *et al.*, *Genomics* 11:508 (1991); Trackman *et al.*, *Biochemistry* 29:4863 (1990)).

Oxidation of lysine residues in tropoelastin leads to the formation of α -aminoadipic- δ -semialdehyde (AAS). Reaction of a lysine residue with AAS leads to the formation of a dehydrolysinonorleucine cross-link. Reaction of two AAS residues leads to an Aldol condensation product. Tetrafunctional cross-linkages can be formed
5 in elastin from three peptidyl aldehydes (AAS) and one unmodified lysine residue (Figure 1) (Kagan, *Path. Res. Pract.* 190:910, 1994; incorporated herein by reference).

For the purposes of the present invention, lysyl oxidase can be produced by any available method, including, for example, chemical synthesis, recombinant
10 methods, and purification from a natural source. In a preferred embodiment, lysyl oxidase is produced recombinantly in much the same manner as the tropoelastin (discussed *supra*). Again, lysyl oxidase is normally secreted so a preferred method of producing and purifying the enzyme would be to secrete it into the media and then purify the protein from the media. If a modified version (*i.e.*, change in amino acid
15 sequence) was needed, the appropriate cDNA could be introduced into the cell to produce the desired mutant protein. Standard techniques for these procedures are known in the art.

In a preferred embodiment of the invention, a modified version of the lysyl oxidase is produced which is inactive initially and can subsequently be converted into
20 an active form. This particular embodiment of the invention allows the tropoelastin and inactive lysyl oxidase to be stored together. Then, prior to application to a wound, the inactive lysyl oxidase is converted to the active form by cleavage of the

protein or by a change in pH, temperature, salt concentration, metal ion concentration, *etc.* The conversion from the inactive form to the active form might even take place at the site of the wound and be caused by a protease or change in pH at the wound site *per se*.

5 In another preferred embodiment of the invention, a modified version of lysyl oxidase would be provided which binds to another protein or peptide (binding peptide or protein). The binding of this binding protein or peptide to the modified lysyl oxidase would lead to inactivation of the lysyl oxidase enzyme. In order for the lysyl oxidase to regain its enzymatic activity, the binding protein or peptide would need to
10 be removed. Examples of this approach include competing off the modified lysyl oxidase using another peptide or protein, a small organic molecule, a metal, nucleic acid, polysaccharide, *etc.*; cleavage of the binding peptide or protein; conformational change of modified lysyl oxidase; and conformational change of the binding protein or peptide.

15 In a particular preferred embodiment, a protein from the family of chaperonins (*e.g.*, GroEL, GroES) or heat shock proteins (*e.g.*, Hsp60, Hsp40, Hsp70) may be used as the binding protein, and the lysyl oxidase may be modified by techniques well known in the art to include a binding domain specific for a chaperonin or heat shock protein (Ranson *et al*, "Chaperonins" *Biochem. J.* 333 (Pt.
20 2):233-242, July 15, 1998; Fink *et al.*, "Chaperone-mediated protein folding" *Physiol. Rev.* 79(2):425-449, April 1999; each of which is incorporated herein by reference). In this way, the virgin tropoelastin monomers and modified lysyl oxidase

bound to a chaperonin or heat shock protein can be stored together without the risk of cross-linking the tropoelastin monomers. After the tropoelastin and lysyl oxidase are applied to a wound or immediately before they are applied to a wound, an agent such as a small molecule (*e.g.*, estrogen) is added to restore lysyl oxidase activity and

5 begin the cross-linking of tropoelastin (Bohen *et al.*, "Hold 'em and fold 'em: chaperones and signal transduction" *Science* 268(5125):1303-1304, June 2, 1995; incorporated herein by reference).

Packaging

10 In one preferred embodiment of the invention, the tropoelastin monomers are be provided in one vial, and the lysyl oxidase are provided in another vial. These proteins might be provided in a dry powder form, in solid form (*i.e.*, lyophilized), in solution, or in suspension. To the proteins may have been added emulsifiers, salts, preservatives, other proteins, nucleic acids, protease inhibitors, antibiotics, perfumes,

15 polysaccharides, adhesive agents, polymers, microfibrils, oils, *etc.*

In another preferred embodiment, the tropoelastin or the lysyl oxidase, or both, is encapsulated in a biodegradable polymer so that when the composition is applied to a wound, the polymer degrades and the tropoelastin and/or lysyl oxidase is released; the released tropoelastin can then be crosslinked.

20 Biodegradable polymers are usually based on functional groups such as esters, anhydrides, orthoesters, and amides. Rapidly biodegradable polymers include poly[lactide-co-glycolide], polyanhydrides, and polyorthoesters. Preferred bioerodible

polymers include polylactides, polyglycolides, and copolymers thereof, poly(ethylene terephthalate), poly(butyric acid), poly(valeric acid), poly(lactide-co-caprolactone), poly(lactide-co-glycolide), polyanhydrides, polyphosphazenes, poly(ϵ -caprolactone), poly(dioxanone), poly(hydroxybutyrate), poly(hydroxyvalerate), polyorthoesters, blends, and copolymers thereof. Examples of biodegradable and biocompatible polymers of acrylic and methacrylic acids or esters include poly(methyl methacrylate), poly(ethyl methacrylate), poly(butyl methacrylate), poly(isobutyl methacrylate), poly(hexyl methacrylate), poly(isodecyl methacrylate), poly(lauryl methacrylate), poly(phenyl methacrylate), poly(methyl acrylate), poly(isopropyl acrylate), poly(isobutyl acrylate), poly(octadecyl acrylate), *etc.* Other polymers which can be used in the present invention include polyalkylenes such as polyethylene and polypropylene; polyarylalkylenes such as polystyrene; poly(alkylene glycols) such as poly(ethylene glycol); poly(alkylene oxides) such as poly(ethylene oxide); and poly(alkylene terephthalates) such as poly(ethylene terephthalate). Additionally, polyvinyl polymers can be used which include polyvinyl alcohols, polyvinyl ethers, polyvinyl esters, and polyvinyl halides. Exemplary polyvinyl polymers include poly(vinyl acetate), polyvinyl phenol, and polyvinylpyrrolidone. Mixtures of two or more of the above polymers could also be used in the present invention.

Some polymeric materials are known to release entrapped compounds upon exposure to a stimulus such as a change in pH or temperature. An examples of microparticles that release as a function of a change in pH include the diketopiperazine particles describes in U.S. Patent 5,352,46,1 issued on October 4,

1994, to Steiner *et al.*, and the proteinoid formulations described in U.S. Patent Reissue no. 35,862, issued on July 28, 1998.

Application

5 Methods for using the composition include, for example, mixing the two separate components together immediately before application to the wound (*i.e.*, before substantial cross-linking has occurred). Other preferred embodiments include applying either the tropoelastin or lysyl oxidase and then the other one to the wound site. One particularly preferred embodiment involves application of a composition
10 comprising lysyl oxidase and tropoelastin to a wound and then use of sutures, staples, adhesive strips, or tissue adhesive to hold the tissue together during the healing process.

 Another particularly preferred embodiment involves application of the composition more than once during the healing process. Preferred regiments for
15 applying the lysyl oxidase and tropoelastin include several times a day, once a day, once a week, twice a week, once a month, and twice a month. In following each of these regiments, the agents would be mixed immediately before application, or they would be applied separately to the wound.

 The lysyl oxidase and tropoelastin are delivered to the wound site using any
20 means to apply a liquid, paste, gel, or solid (*e.g.*, powder). These means include, for example, a brush, a syringe, a spatula, and a container specifically designed to delivery the agents such as a tube with a narrow tip.

In another preferred embodiment of the invention, the composition is applied to a wound resulting from deep trauma or surgery involving the lungs, large arteries, or other tissues with significant amounts of elastic fibers.

5

Other Embodiments

The foregoing has been a description of certain non-limiting preferred embodiments of the invention. Those of ordinary skill in the art will appreciate that various changes and modifications to this description may be made without departing from the spirit or scope of the present invention, as defined in the following claims.

10

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000

Claims

What is claimed is:

- 5 1. A method for promoting wound healing comprising the steps of:
- providing isolate tropoelastin and isolated lysyl oxidase
- separated from each other; and
- applying both said tropoelastin and said lysyl oxidase to wound
- simultaneously or sequentially.
- 10 2. The method of claim 1 wherein tropoelastin is wild type
- tropoelastin matched to species of recipient.
3. The method of claim 1 wherein tropoelastin is modified
- 15 tropoelastin.
4. The method of claim 3 wherein amino acid sequence of
- tropoelastin has been changed relative to amino acid sequence of wild
- type tropoelastin.
- 20 5. The method of claim 1 wherein tropoelastin comprises a heterogeneous
- mixture of tropoelastin isoforms.

6. The method of claim 1 wherein lysyl oxidase comprises an enzymatically active portion of lysyl oxidase.
7. The method of claim 1 wherein lysyl oxidase is modified lysyl oxidase relative to wild type lysyl oxidase.
8. The method of claim 7 wherein the amino acid sequence of lysyl oxidase has been changed relative to wild type lysyl oxidase.
9. The method of claim 1 wherein the method comprises the additional step of:
- repeatedly applying the tropoelastin and lysyl oxidase to the wound during the healing process.
10. The method of claim 1 wherein the method comprises the additional step of:
- approximating separated tissue of the wound using sutures, staples, adhesive strips, or tissue glue.
11. The method of claim 1 wherein the step of applying comprises applying the tropoelastin and lysyl oxidase with a sterile syringe.

12. The method of claims 1 wherein the tropoelastin or lysyl oxidase has been mixed with other materials selected from the group consisting of polymers, emulsifiers, oils, perfumes, proteins, polysaccharides, nucleic acids, microfibrils, antimicrobial agents, adhesive agents, and protease inhibitors.
- 5
13. A kit comprising tropoelastin and lysyl oxidase in separate compartments.
- 10
14. The kit of claim 13 wherein the tropoelastin is wild type tropoelastin.
15. The kit of claim 13 wherein the tropoelastin is modified tropoelastin.
- 15
16. The kit of claim 15 wherein the amino acid sequence of tropoelastin has been changed relative to wild type sequence.
17. The kit of claim 13 wherein the lysyl oxidase is wild type lysyl oxidase.
- 20
18. The kit of claim 13 wherein the lysyl oxidase is modified lysyl oxidase.

19. The kit of claim 18 wherein the amino acid sequence of lysyl oxidase has been changed relative to wild type sequence.

5 20. A kit comprising tropoelastin and lysyl oxidase in the same compartment, wherein the lysyl oxidase is in an inactive form which can be later converted to an active form.

10 21. A kit comprising tropoelastin and lysyl oxidase in the same compartment, wherein either the tropoelastin or lysyl oxidase is encapsulated by a polymer.

22. The kit of claim 21 wherein the polymer is biodegradable.

15 23. The method of claim 1 wherein the wound involves the skin.


24. The method of claim 1 wherein the wound involves an artery.

25. The method of claim 1 wherein the wound involves lung tissue.



Abstract

The present invention provides compositions and methods for promoting wound healing. The composition comprises virgin monomers of tropoelastin and lysyl oxidase. When the lysyl oxidase comes in contact with the tropoelastin, cross-
5 linking of the tropoelastin monomers will occur to form elastin. Contacting the tropoelastin and lysyl oxidase together and applying the mixture to a wound before substantial cross-linking has occurred promotes wound healing by holding the damaged tissue together, increasing the rate of healing, and decreasing the amount of scarring.

10



Non-enzymatic Condensation Cross-Linking Reactions



DECLARATION

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name,

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

WOUND HEALING COMPOSITIONS AND METHODS USING TROPOELASTIN AND LYSYL OXIDASE

the specification of which (I authorize Choate, Hall & Stewart to check one of the following, three choices, and fill in the blanks, if applicable):

 X is attached hereto

 was filed on as Application
Serial No. and amended on (if applicable).

 was filed as PCT international application No. ,
on and was amended under PCT Article 19
on (if applicable).

I hereby state that I have reviewed and understood the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

I acknowledged the duty to disclose information which is material to the examination of this application in accordance with Title 37, Code of Federal Regulations, §1.56.

I hereby claim foreign priority benefits under Title 35, United States Code, §119 of any foreign application(s) for patent or inventor's certificate listed below and have also identified below any foreign application for patent or inventor's certificate having a filing date before that of the application on which priority is claimed:

Prior Foreign Application(s):

Priority Claimed

<u> </u> (Number)	<u> </u> (Country)	<u> </u> (Day/Month/Year/Filed)	<u> </u> Yes	<u> </u> No
-------------------------------	--------------------------------	---	----------------------	---------------------

<u> </u> (Number)	<u> </u> (Country)	<u> </u> (Day/Month/Year/Filed)	<u> </u> Yes	<u> </u> No
-------------------------------	--------------------------------	---	----------------------	---------------------

I hereby claim the benefit under Title 35, United States Code, §120 of any United States application(s) or PCT international application(s) designating the United States of America listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code, §112, I acknowledge the duty to disclose material information as defined in Title 37, Code of Federal Regulations, §1.56 which became available between the filing date of the prior application and the national or PCT international filing date of this application:

_____ (Application Serial No.)	_____ (filing date)	_____ (status-patented, pending, abandoned)
-----------------------------------	------------------------	--

_____ (Application Serial No.)	_____ (filing date)	_____ (status-patented, pending, abandoned)
-----------------------------------	------------------------	--

PCT Applications designating the United States:

_____ (PCT Appl. No.)	_____ (U.S.S.N.)	_____ (status-patented, pending, abandoned)
--------------------------	---------------------	--

I hereby claim the benefit under Title 35, United States Code, §119(e) of any United States provisional application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code, §112, I acknowledge the duty to disclose information which is material to patentability as defined in Title 37, Code of Federal Regulations, §1.56 which became available between the filing date of the prior application and the national filing date of this application.

Provisional Application(s):

_____ Application Number	_____ Filing Date	_____ Status
-----------------------------	----------------------	-----------------

_____ Application Number	_____ Filing Date	_____ Status
-----------------------------	----------------------	-----------------

_____ Application Number	_____ Filing Date	_____ Status
-----------------------------	----------------------	-----------------

[illegible]

ATTORNEY'S DOCKET NO.: 2001605-0007

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicant: Ensley, Burt D.
Serial Number:
Filed: February 4, 2000
For: WOUND HEALING COMPOSITIONS AND METHODS USING
TROPOELASTIN AND LYSYL OXIDASE

Assistant Commissioner for Patents
Washington, D.C. 20231

Sir:

APPOINTMENT OF ATTORNEY

The undersigned hereby appoints Brenda Herschbach Jarrell, Registration No. 39,223; Sam Pasternack, Registration No. 29,576; Kevin M. Tormey, Registration No. 41,351; Elizabeth Nugent, Registration No. 43,839, Valarie B. Rosen, Registration No. P-45,698; Stanley C. Mah, Registration No. P-46,189 and Karoline M. Shair, Registration No. 44,332 as its attorneys and agents for prosecution of matters relating to the above-identified patent application and to conduct all business in the United States Patent and Trademark Office.

All correspondence should be sent to Brenda Herschbach Jarrell, Choate, Hall & Stewart, Exchange Place, 53 State Street, Boston, Massachusetts 02109.

Respectfully submitted,

Name: _____
Title: _____
On behalf of _____

Dated: _____, 2000